ISOLATION AND CHARACTERIZATION OF COW DUNG MICROFLORA AND ITS EFFECTIVENESS OF *CITRUS LIMON* BIO-ENZYME

Abstract

Cow manure act as a good fertilizer. It contain more nutrients beneficial microbes. It supports growth useful microbes. When it was mixed with soil it improve the soil and maintain moisture. The present studies carried out isolation of cow dung microflora and its antimicrobial properties against bio enzyme *citrus limon*. The bacterium like *shigella* species *23mm* inhibition and proteous species17 mm inhibition against *citrus limon*.

Authors

Krishnaveni R

Assistant professor and Head Department of Microbiology Idhaya College for Women Kumbakonam, Tamil Nadu, India krishnavenimicro@gmail.com

Sathya I

UG Student Department of Microbiology Idhaya College for Women Kumbakonam, Tamil Nadu, India

Ameera S

UG Student Department of Microbiology Idhaya College for Women Kumbakonam, Tamil Nadu, India

Arthi K

UG Student Department of Microbiology Idhaya College for Women Kumbakonam, Tamil Nadu, India

Janaranjani S

UG Student Department of Microbiology Idhaya College for Women Kumbakonam, Tamil Nadu, India

I. INTRODUCTION


Cow dung more contain more beneficial microbes and various Nutrional components vitamins, cellulose, oxygen, carbon, mucus, potassium, nitrogen, it is being used in Agricultural and religious purpose.

A micro organisms such as *bacilli coccus*. According to *Ware et al.*, 1988 Lactation performance of two large dairy herds fed Lactobacillus acidophilus strain. Saccharomyces corevisiae act as probiotic Ware Funssin D R, Read PL et AL., (1988) Generally old cow dung has more soil microbes and Actinomices Muhamed and Amusha (2003). There are many Proofs to confirm that. Nene YL. (2001) Utilizing traditional knowledge in agriculture. Traditional knowledge system of India and Sri Lanka. Sharma and Singh, 2015)). Isolation and charactreization of bacteria from cow dung of desi cow breed on different morpho-biochemical parameters in Dehradun. (Randhawa and Kullar, 2011) Bioremediation of pharmaceuticals, pesticides and petrochemicals with gomeya dung. (Fleming et.al., 1929) Kardos N and Demain AL. (2011). Cow dung and cow rich in Enterobacter aerogenes, Escherichia coil, Klebsilla oxytoca, Klebsilla pneumonia Morgarella morganii, Pasteurella species, antifungal agent. (Muhamed and Amusha 2003) (Dhama et al., 2005). (Joseph and Sankerganesh 2001) (Dhama et al., 2013) cow dung shows antimicrobial properties (Daviud odemi et al., 2007)

Microbial Analysis of Compost Using Cow Dung as Booster. Cow dung as booster in the decomposition of organic material (*Adgunloye et al., 2007*). Cow dung act as organic Fertilizer, (*V. Muralikrishna et al., 2017*) Recycling of Organic Wastes in Agriculture: (*B. Sharma et al., 2019*)



Citrus limon HerbalPlant



Bio Enzyme Citruslimon.,

Bio Chemical Charecteristic Of Shigella Species.,


Futuristic Trends in Biotechnology ISBN: 978-93-95632-84-3 IIP Proceedings, Volume 2, Book 28, Part 1, Chapter 16 ISOLATION AND CHARACTERIZATION OF COW DUNG MICROFLORA AND ITS EFFECTIVENESS OF CITRUS LIMON BIO-ENZYME




Shigella Sp, Hekteon Entric Agar




MR -Positive



Urease Test-Positive




Indole NegativeTest



Indole Negative



Klebisella Pneumoniae Species



Simmon's Citrate Test- Positive



Biochemical characters of *Proteus species*

Simmon's Citrate Test - Positive

 Table: 4 Testing Antimicrobial Sensitivity of Isolates Against Bio – Enzyme (Citrus Limon)

sl.no	Tested Organism	Zone Of Inhibition
1.	Klebsiella species	14mm
2.	Proteus species	15mm
3.	Shigella species	17mm

Antimicrobial activity of isolated microbes



Klebisella Pneumoniae



Disc diffusion method- Shigella Species.,



Disc diffusion method- Proteus species.,

II. RESULT AND DISSCUSSION

The microbes like *Klebsiella* species, *shigella* species, and *Proteus* species isolated from cow dung. Testing of antimicrobial activity of isolates with *Citrus limon Klebsiella species* 14 mm in zone of inhibition in muller hinton Agar medium. *Proteus* species 15 mm in zone of inhibition, *Shigella* species 17 mm zone of inhibition observed.this study concluded Citrus peels bio enzyme is a antimicrobial agent. The present work concluded that the bioenzyme of *Citrus limon* has antimicrobial properties .it ia used as cleanising solution.

III.REFERENCES

- [1] Abo-State MA, Mahdy HM, Ezzat SM, Abd El Shakour EH and ElBahnasawy MA. (2012). Antimicrobial resistance profiles of Enterobacteriaceae isolated from rosettabranchofriverNile,Egypt.WorldAppliedSciencesJournal.19:1234-1243.
- [2] Adegunloye DV, Adetuyi FC, Akinyosoye FA, Doyeni MO (2007). Microbial Analysis of Compost Using Cow Dung as Booster.Pak.J.Nutri.6(5):506-510.
- [3] David OM, Odeyemi AT (2007). Antibiotic resistant pattern of environmental isolates of Listeria monocytogenes from Ado-Ekiti, Nigeria. Afr. J. Biotechnol. 6(18):2135-2139
- [4] Dhama K, Chakraborty S and Tiwari R. (2013). Panchgavya therapy (Cowpathy) in safeguarding health of animals and humans-a review. Res OpinAnim Vet Sci. 3:170-178.
- [5] Fleming A. (1929). On the antibacterial action of cultures of a penicillium, with special reference to their use in the isolation of B. influenzae. Br. J. Exp. Pathol. 10:226–236.
- [6] Joseph B and Sankarganesh P. (2011). Antifungal efficacy of panchgavya.
- [7] International Journal of PharmTech Research. 3:585-588.
- [8] Kardos N and Demain AL. (2011). Penicillin: the medicine with the greatest impact on therapeutic outcomes. Appl. Microbiol. Biotechnol. 92:677–687.
- [9] Muralikrishna.V., V. Manickam, I. V. Muralikrishna, V. Manickam, Solid
- [10] Waste Management, Environ. Manage. (2017) 431-462. https://doi.org/10.1016
- [11] /B9 78-0-12-811989-1.00016-6.
- [12] Randhawa GK, Kullar JS (2011) Bioremediation of pharmaceuticals, pesticides,
- [13] and petrochemicals with gomeya/cow dung. ISRN Pharmacol.
- [14] doi:10.5402/2011/362459
- [15] Sharma, B. and Singh, M. (2015). Isolation and charactreization of bacteria from cow dung of desi cow breed on different morpho-biochemical parameters in Dehradun. International journal of Advances in Pharmacy, Biology and Chemistry, Vol:4(2).
- [16] B. Sharma, B. Vaish, Monika, U.K. Singh, P. Singh, R.P. Singh, Recycling of Organic Wastes in Agriculture: An Environmental Perspective, Int. J. Environ. Res. 13 (2019) 409– 429.https://doi.org/10.1007/s41742-019-00175-y
- [17] Ware Fungsin, D.R., Read, P.L., Mantredi, E.T. (1988). Lactation performance of two large dairy herds fed Lactobacillus acidophilus strain BT 1386. J. Dairy Sci, 71:219-222.